

# TWO-LOOP RENORMALIZATION OF THE QUASIPARTICLE WEIGHT IN TWO-DIMENSIONAL ELECTRON SYSTEMS

Jun-ichiro Kishine

*Department of Theoretical Studies, Institute for Molecular Science, Okazaki 444-8585, Japan*

*E-mail: kishine@ims.ac.jp*

Nobuo Furukawa

*Department of Physics, Aoyama Gakuin University, Setagaya, Tokyo 157-8572, Japan*

*E-mail: furukawa@phys.aoyama.ac.jp*

Kenji Yonemitsu

*Department of Theoretical Studies, Institute for Molecular Science, Okazaki 444-8585, Japan*

*E-mail: kry@ims.ac.jp*

We apply the renormalization-group (RG) approach to two model systems where the two-dimensional Fermi surface has portions which give rise to the logarithmically singular two-loop self-energy process.

## 1 Introduction

Non-Fermi-liquid behavior in the normal state of high  $T_c$  cuprates has driven us to renewed interest in understanding the two-dimensional (2D) interacting electron systems in the framework of the renormalization-group (RG) approach.<sup>1</sup> In some cases, a 2D Fermi surface has portions which give rise to coupled infrared singularities in the particle-particle and particle-hole fluctuations. Many attempts have so far been made at detecting instabilities of the Landau-Fermi-Liquid by keeping track of the *one-loop* RG flow of the corresponding susceptibilities. Among them are included the 2D systems with a partially flat Fermi surface,<sup>2,3</sup> tight-binding dispersion,<sup>4,5</sup> and van-Hove singularity.<sup>6,7</sup> The breakdown of the Landau-Fermi-liquid is also signaled by the vanishing quasiparticle weight,  $z$ , which is treated on the *two-loop* level. In this paper, we apply the two-loop RG analysis to two model systems where the 2D Fermi surface has portions which give rise to the infrared logarithmic singularity (ILS) of the two-loop self-energy (2LSE).

## 2 Fermi surface with partially flat regions

As the first example, we consider a 2D Fermi surface shown in Fig. 1(a), which consists of flat regions ( $\alpha, \bar{\alpha}, \beta, \bar{\beta}$ ) with length  $2\Lambda$  and round-arc regions.<sup>2,3</sup> We linearize the one-particle dispersion with the bandwidth  $E_0$  in the flat regions and parametrize the infrared cutoff as  $\omega_l = E_0 e^{-l}$  with a scaling parameter,  $l$ . We assume  $E_0 \ll v_F \Lambda$  with  $v_F$  being the corresponding Fermi velocity. In this case, the RG equations are derived as in one dimension except that there appear infinite numbers of two-particle scattering vertices connecting the parallel flat regions [Fig. 1(b)].<sup>2</sup>

Figure 1: (a) The Fermi surface considered here. (b) The two-particle scattering vertices connecting the flat regions  $\alpha$  and  $\bar{\alpha}$ , where  $-\Lambda \leq k_{y1}, k_{y2}, k_{y3} \leq \Lambda$  and  $\sigma_1, \dots, \sigma_4$  are spin indices. (c) The two-loop self-energy diagram. The solid and broken lines represent the propagators for the electrons in the flat regions,  $\alpha$  and  $\bar{\alpha}$ , respectively.

Figure 2:  $k_y$  dependence of  $z_l(k_y)$  at each RG step.

The RG equation for the quasiparticle weight at the position  $k_y$  in the region  $\alpha$ ,  $z(k_y)$ , is written as<sup>3</sup>

$$d \ln z(k_y) / dl = -\theta_l(k_y), \quad (1)$$

where  $\theta_l(k_y)$  comes from the 2LSE diagram shown in Fig. 2(c) and contains the scattering vertices which are also treated on the two-loop level. As an initial condition at  $l = 0$ , we put the quasiparticle weight equal to the non-interacting value,  $z_0(k_y) = 1$ , and the scattering strengths corresponding to the vertices in Fig. 1(b) equal to the Hubbard repulsion,  $U$ .

In Fig. 2, we show the RG flow of  $z(k_y)$  for  $U = 4.5\pi v_F$ . We see that the region around which  $z_l(k_y)$  is the most strongly suppressed moves from the center ( $k_y = 0$ ) for  $l < l_{\text{cross}} \sim 3$  toward the edges ( $k_y = \pm\Lambda$ ) for  $l > l_{\text{cross}}$  as the energy scale decreases and finally  $z_l(k_y)$  approaches zero everywhere for  $-\Lambda \leq k_y \leq \Lambda$  in the low-energy limit. The crossover behavior originates from the breakdown of the cancellation between the particle-particle and particle-hole loops.<sup>3</sup>

### 3 Fermi surface which touches the umklapp surface

As the second example, we consider a 2D Fermi surface shown in Fig. 3(a), which touches the umklapp surface at 4 points,  $(\pm\pi/2, \pm\pi/2)$ .<sup>8,9</sup> We divide the Fermi surface into patches and take account of the electron one-particle and two-particle processes within the patches  $\alpha = 1, 2, 3, 4$  around  $(\pm\pi/2, \pm\pi/2)$  points [see Fig. 3(a)] which are connected through the normal  $[g_1, g_2, g_{1r}]$  and the umklapp  $[g_3, g_{3p}, g_{3x}]$  processes,<sup>8,9</sup> as indicated in Fig. 3(b). We linearize the one-particle dispersion with

Figure 3: (a) The Fermi surface considered here. (b) The two-particle scattering vertices entering the RG equations. (c) The two-loop self-energy diagram. Each leg of the vertices contains a patch index  $\alpha = 1, 2, 3, 4$ .

the bandwidth  $E_0$  in the direction normal to the Fermi surface at the center of the patch and parametrize the infrared cutoff as  $\omega_l = E_0 e^{-l}$  with a scaling parameter,  $l$ . In the energy scale,  $\omega > \omega_T$ , where  $\omega_T$  is the infrared cutoff from the transverse dispersion, the RG equations are derived as in one dimension except that there appear additional inter-patch vertices,  $g_{1r}$ ,  $g_{3p}$ , and  $g_{3x}$ .

The 2LSE diagram, shown in Fig. 3(c), gives rise to the ILS for particular pairs of patch indices,  $(\alpha, \beta) = (1, 3)$  and  $(2, 4)$ . The RG equation for the quasiparticle weight at each patch,  $z$ , is

$$d \ln z / dl = -(g_1^2 + g_2^2 - g_1 g_2 + g_3^2 / 2) / 4. \quad (2)$$

The two-particle scattering vertices are also treated on the two-loop level:

$$dg_1 / dl = -g_1^2 - g_{1r}^2 - 2g_{3x}^2 + 2g_{3x}g_{3p} - g_1^3 / 2, \quad (3)$$

$$dg_2 / dl = -(g_1^2 + 2g_{1r}^2 - g_3^2 - 2g_{3p}^2) / 2 - g_1^3 / 4 + g_3^2(g_1 - 2g_2) / 4, \quad (4)$$

$$dg_{1r} / dl = -(g_1 + g_2)g_{1r}, \quad (5)$$

$$dg_3 / dl = -(g_1 - 2g_2)g_3 - 2g_{3x}^2 + 2g_{3x}g_{3p} + g_{3p}^2 - g_3(g_1 - 2g_2)^2 / 4 - g_3^3 / 4, \quad (6)$$

$$dg_{3x} / dl = -2g_1g_{3x} + g_1g_{3p} + g_2g_{3x} - g_3g_{3x} + g_3g_{3p}, \quad (7)$$

$$dg_{3p} / dl = (g_2 + g_3)g_{3p}. \quad (8)$$

The RG equations for  $g_{1r}$ ,  $g_{3p}$ , and  $g_{3x}$  processes do not contain two-loop contribution, on the same ground that logarithmically singular contribution of the 2LSE is limited to particular pairs of patch indices. As an initial condition at  $l = 0$ , we put the quasiparticle weight equal to the non-interacting value,  $z_0 = 1$ , and all the scattering strength corresponding to the vertices in Fig. 3(b) equal to the Hubbard repulsion,  $U$ .

In Fig. 4(a), we show the two-loop RG flow of the scattering strengths for  $U = 0.5\pi v_F$ . As in the case of the one-loop analysis,<sup>9</sup> all the umklapp scatterings,  $g_3$ ,  $g_{3p}$ , and  $g_{3x}$ , are renormalized to a strong coupling sector for  $U > 0$ . By examining the spin and charge susceptibilities, Furukawa and Rice suggested that the strong coupling fixed point corresponds to an insulating spin liquid state at the 4 patches around  $(\pm\pi/2, \pm\pi/2)$ .

In Fig. 4(b) are shown the RG flows of the quasiparticle weight at the 4 patches,  $z$ , in the following two cases: **(i)**  $g_1$ ,  $g_2$ , and  $g_3$  are present and other vertices

Figure 4: (a) The two-loop RG flow of the scattering strengths for  $U = 0.5\pi v_F$ . (b) The RG flows of  $z$  in the cases where (i)  $g_1$ ,  $g_2$ , and  $g_3$  are present and other vertices are absent, and (ii) all vertices shown in Fig. 3(b) are present.


are absent, and (ii) all the vertices shown in Fig. 3(b) are present. The case (i) is equivalent to the case of one dimension. We see that the  $g_{1r}$ ,  $g_{3p}$ , and  $g_{3x}$  processes, which are peculiar to the 2D Fermi surface considered here, strongly suppress the quasiparticle weight. Consequently the quasiparticle weight at the 4 patches approaches zero at a energy scale much higher than in the case (i). This result complements the view based on the one-loop analysis.<sup>9</sup>

In both the first and second examples taken up here, we have neglected effects of a Fermi surface curvature on the RG equations. If we take the finite curvature into account, the RG flow of the quasiparticle weight might stop around the energy scale  $\omega_{\text{curv}}$  corresponding to the resolution which detects the curvature. It remains matter for debate how to incorporate the energy scale  $\omega_{\text{curv}}$  into the RG-based scheme in 2D.


## Acknowledgments

J. K thanks J. B. Marston for helpful comments during the MBX conference. N. F. thanks T. M. Rice and M. Salmhofer for discussion. This work was supported by a Grant-in-Aid for Encouragement of Young Scientists from the Ministry of Education, Science, Sports and Culture, Japan.


1. R. Shankar, Rev. Mod. Phys. **66**, 129 (1994).
2. A. T. Zheleznyak V. M. Yakovenko, and I. E. Dzyaloshinskii, Phys. Rev. **B 55**, 3200 (1997).
3. J. Kishine and K. Yonemitsu, Phys. Rev. **B 59**, 14823 (1999).
4. D. Zanchi and H. J. Schulz, Phys. Rev. **B 54**, 9509 (1996).
5. C. J. Halboth and W. Metzner, preprint, cond-mat/9908471.
6. H. J. Schulz, Europhys. Lett. **4**, 609 (1987).
7. N. Furukawa, T. M. Rice, and M. Salmhofer, Phys. Rev. Lett. **81**, 3195 (1998).
8. A. Houghton and J. B. Marston, Phys. Rev. **B 48**, 7790 (1993).
9. N. Furukawa and T. M. Rice, J. Phys. Cond. Mat. **10**, L381 (1998).




Kishine, Furukawa, and Yonemitsu Fig. 1



Kishine, Furukawa, and Yonemitsu Fig. 2



Kishine, Furukawa, and Yonemitsu Fig. 3



Kishine, Furukawa, and Yonemitsu Fig. 4